

Innovative Strategies to Pelvic Trauma Pain Management: A Systematic Narrative Review of Pharmacological, Regional, and Multidisciplinary Advances

Dr. Anushya¹, Dr. R. R. Redlin Jani², S. Grace³, T. Jayashree⁴, B. S. Janani⁵, M. Gayathri⁶ and M. Devi Varshini⁷

¹Physician, Department of Emergency, Vivekanandha Medical Care Hospital, INDIA

²Assistant Professor, Department of Pharmacy Practice, Swamy Vivekanandha College of Pharmacy, INDIA

^{3,4,5,6,7}V Pharm D, Department of Pharmacy Practice, Swamy Vivekanandha College of Pharmacy, INDIA

³Corresponding Author: grace07pharmd2020@gmail.com

ORCID

<https://orcid.org/0009-0002-9218-5083>



www.jrasb.com || Vol. 4 No. 4 (2025): August Issue

Received: 09-08-2025

Revised: 13-08-2025

Accepted: 18-08-2025

ABSTRACT

The unique anatomy and associated effects of pelvic trauma, which frequently arises from high-energy traumatic events like falls or vehicle collisions, make pain management extremely challenging. Although its efficiency, traditional opioid-based analgesia has disadvantages like tolerance, reliance, and respiratory impairment. With emphasis on pharmacologic advancements, regional anesthesia, minimally invasive techniques, and multidisciplinary approaches, this systematic review illustrates recent developments in pelvic trauma pain management. Non-opioid medicines which include gabapentinoids, acetaminophen, and NSAIDs are becoming significant components of multimodal analgesia. While strengthening pain management, adjuncts such as intravenous lidocaine and ketamine reduce the usage of opioids. For prolonged local analgesia, liposomal bupivacaine along with additional targeted methods of administration are available. Sacral nerve blocks, fascia iliaca, and lumbar plexus are a few regional treatments that successfully alleviate pain at specific sites. In instances where medication fails to benefit the patient, minimally-invasive techniques including spinal cord stimulation (SCS), peripheral nerve stimulation, and radiofrequency ablation (RFA) show promise. Nowadays, the standard of excellence is an integrated approach which involves pharmacologic, interventional, psychological, and rehabilitative ways. nevertheless, challenges which includes insufficient funding, disparate procedures, and insufficient guidance prevent broad adoption. To enhance the results of pelvic trauma care, future initiatives should give special focus to standardization, long-term evaluation, and equal access to these innovative therapies.

Keywords- Pelvic trauma, Multimodal pain management, Regional anesthesia, Opioid-sparing strategies, Minimally invasive techniques.

I. INTRODUCTION

High-energy incidents such as falls or car accidents often result in pelvic trauma, which is defined as injuries to the pelvic bones, joints, and soft tissues. Complicated fracture patterns, excessive bleeding, and significant damage to organs nearby make these injuries problematic. The genital system, the bladder, and major

blood veins are among the most critical organs positioned in the pelvis, causing these injuries possibly fatal as well as difficult to repair ^[1]. The Young-Burgess system classifies pelvic fractures depending on the mechanism of injury: anteroposterior compression (APC) leads to "open book" injuries; lateral compression (LC) causes parallel pubic rami fractures and sacral compression; vertical shear (VS) results in superior hemipelvic displacement;

and multiple force vectors are involved in combined mechanism (CM) fractures. With the goal to offer guidance on treatment and prognosis, these companies assess pelvic stability along with prospective outcomes^[1]. When the bleeding affects the presacral venous plexus or arterial branches, it is one of the primary causes of death. Additional risks include genitourinary injuries, such as urethral dislocation or bladder rupture. While surgical alternatives have been used in the past, new surgical techniques such as percutaneous stabilization have generated superior outcomes (Cothren et al., 2007)^[2].

These developments aside, pelvic trauma has been correlated with a higher risk of morbidity and mortality. The research of Huitinen et al. (2022), an integrated approach is essential to enhancing patient recovery from complex fractures and bleeding^[3]. Pelvic trauma, involving pelvic ring fractures along with associated soft tissue and visceral injuries, remains an extremely unusual yet overwhelmingly deadly category of trauma. An average incidence of 34.3 per 100,000 people was determined by a statewide U.S. database investigation. The hospital mortality rate was 7%, and the case mix of elderly, comorbidities patients was incrementally increasing. A disability adjusted life years (DALYs) have grown by 39% since 1990, according to global data from the 2021 World Burden of Disease project. This rise in disability is being caused by older populations and prolonged contact to high energy methods^[4]. Because of the biomechanical complexity of the pelvic ring, the impact's vector and magnitude influence the patterns of injury. Each pattern carries a unique instability profile and bleeding risk. The Young-Burgess classification divides lesions into four distinct groups: anteroposterior compression, lateral compression, vertical shear, and mixed mechanism types. Long-term prognostication, definitive fixation methodologies, and resuscitation priorities are thus guided by accurate early classification^[5]. Even with contemporary resuscitation methodologies, up to one-third of premature deaths are still caused by exsanguinating bleeding from ruptured iliac branch arteries or pelvic venous plexuses, and in many large volume hospitals, total fatalities from cases still surpass 10%. Thus, it is vital to have coordinated orthopedic, general surgery, and interventional radiology input as well as rapid volume control (pelvic binders, preperitoneal packing, REBOA)^[5].

Pelvic trauma pain is usually severe, multimodal (nociceptive, neuropathic, visceral), and prognostically significant because unmanaged pain hinders the body's ability to heal and recover. A paradigm change toward multimodal analgesia has been prompted by expanding evidence of opioid-related respiratory impairment, delirium, and dependence, which was previously virtually exclusively employed with systemic opioids^[6]. Modern pharmaceutical medications incorporate gabapentinoids, small amounts of ketamine infusions, and non-opioid drugs (NSAIDs, intravenous acetaminophen), the latter of which delivers NMDA-mediated antinociception with

significant opioid sparing effects and hemodynamic neutrality. Low-dose ketamine was shown to be approximately as effective as morphine for treating acute pain in a 2023 systematic analysis of 15 RCTs, with less adverse respiratory and gastrointestinal consequences^[7].

At the same time, catheter technology and ultrasound guidance have enhanced regional anesthetic techniques, such as continuous catheter infusions, sacral plexus or pudendal blocks, and lumbar or fascia iliaca nerve blocks. High degree of confidence Based on Cochrane investigation, peripheral nerve blocks minimize the probability of acute confusional states following hip and pelvic fractures in half and decrease dynamic pain scores by about 2.5/10. These techniques right now serve as a foundation of improved recovery paths for patients with pelvic fractures, together with minimally invasive treatment, early mobilization, and psychological assistance. This article summarizes the current evidence on pharmacologic, regional, and interventional pain management strategies for pelvic trauma against the backdrop of changing epidemiology and therapeutics. It also evaluates implementation barriers in settings with various assets and outlines future research priorities, such as standardizing protocols, establishing advanced drug delivery systems, and using robotic or image-guided minimally invasive techniques, in order to achieve maximum functional recovery and lower mortality^[8].

II. PHARMACOLOGICAL ADVANCEMENTS

Acute pelvic trauma management for pain still heavily depends on traditional analgesics like opioids, however multimodal pharmacological approaches have emerged in response to their adverse effects and the possibility of dependence. To target several pain pathways and lower opioid dosages, they mix opioids with non-opioid analgesics such NSAIDs, Acetaminophen, and adjuvant medications like Gabapentinoids or ketamine. Ahmadi and associates, 2016. In particular in trauma patients facing excruciating pain, ketamine, an NMDA receptor antagonist, has gained popularity for its analgesic and opioid-sparing properties. Low-dose ketamine infusions are ideal for individuals who are hemodynamically unstable because they can effectively relieve pain without causing respiratory depression. Although there is currently little clinical data, new research also investigates more recent drugs that target neuropathic pain mechanisms frequently detected in pelvic injuries, such as selective sodium channel blockers and calcium channel blockers^[9].

1. Non-Opioid Analgesics

Conventional and Innovative Non-Opioid Substances Non-opioid analgesics, which combine both well-known medications and ground-breaking discoveries, have greatly changed as the mainstay of pelvic trauma pain management:

Compared to oral preparations, intravenous NSAIDs and acetaminophen have a quicker onset and better absorption. Faster pain management is essential in trauma situations, and IV ibuprofen produces plasma levels twice as high as those of oral delivery^{[10][11]}. By cutting postoperative opioid use by 20–60%, these formulations greatly reduce the hazards of sedation, nausea, and respiratory depression, supporting opioid-sparing approaches^{[10][12]}. Gabapentinoids, such as gabapentin and pregabalin, work by modulating calcium channels to address neuropathic pain components that frequently occur in pelvic trauma. Pregabalin may offer more quickly and substantial analgesic relief than gabapentin, based on recent systematic reviews; nonetheless, both drugs need cautious dose titration because of possible adverse effects, such as drowsiness and dizziness. For chronic pelvic pain, meta-analyses demonstrate that gabapentin is more effective than a placebo, resulting in significant improvements in pain and mood scores^{[13][14]}. The latest development in non-opioid pain relief is represented by revolutionary novel agents. The first-in-class sodium channel blocker for moderate to severe acute pain, suzetrigine (Journavx), was just approved by the FDA^{[15][16]}. Offering a radical novel approach to acute trauma pain, this mechanism-specific treatment addresses peripheral pain-signaling pathways before the signals reaching the brain^[17]. Selective sodium channel blockers for neuropathic and acute pain syndromes, TRPV1 antagonists that target heat and acid-triggered pain pathways, and nerve growth factor (NGF) monoclonal antibodies for inflammatory pain are among the emerging drug classes being studied^[18].

2. Opioid-Sparing Protocols

Modern opioid-sparing procedures employ systematic, evidence-based methods that integrate several treatment modalities as part of their multimodal analgesia strategies. In terms of trauma pain management, the MAST (Multi-modal Analgesic Strategies in Trauma) Protocol is a significant breakthrough. This generally accessible method consists of: 200 mg of lidocaine every 12 hours; 500 mg of naproxen every 12 hours; 300 mg of gabapentin every 8 hours; 1g of paracetamol PO every 6 hours; and pure opioid formulations only when necessary for breakthrough pain^{[19][20][21]}.

Clinical evidence from randomized controlled studies shows that, in comparison to conventional methods, the MAST protocol maintains equal pain control, lowers opioid prescribing at discharge (62% vs. 67%), and reduces daily opioid intake (34 vs. 48 MME/day)^[22]. This indicates significant opioid decrease without sacrificing analgesic efficacy in trauma patients with median injury severity scores of 14^[20].

In accordance with organized standards, the Stepwise Protocol Implementation begins with regional anesthetic and non-opioid medications, saving opioids for rescue therapy. This method can improve pain scores and functional results while reducing opioid prescriptions by up to 31%^{[23][24][25]}. For improved opioid-sparing effects,

especially in perioperative and critical care settings, Advanced Adjuvant Integration combines specialized medications such as ketamine, dexmedetomidine, and several antidepressants^{[23][26]}.

3. Targeted Drug Delivery Systems

High-Tech Delivery Systems by improving site-specific efficacy and reducing systemic side effects, targeted medication delivery methods transform pain management with implantable pumps, Intrathecal Drug Delivery Systems (ITDD) carry drugs straight to the spinal cord, resulting in therapeutic CNS concentrations with a notably reduced systemic exposure. During the course of 6-29 months of treatment, the aggregate baseline pain scores decreased from 8.70 to 4.45, indicating that these systems are superior at controlling chronic non-malignant pain. About 44.5% of patients report at least 50% less discomfort^{[27][28][29][30]}.

Analgesics are encapsulated in lipid vesicles for targeted distribution and prolonged release according to liposomal drug delivery. The FDA-approved nanosystem for prolonged analgesia, liposomal bupivacaine (Exparel), offers up to 72 hours of pain treatment with no systemic effects^{[31][32][33]}. These sophisticated formulations include PEGylated liposomes with longer circulation times and lower immunogenicity, photo-triggerable liposomes that allow controlled drug release through near-infrared light activation, and stimuli-responsive systems that release medications in response to changes in pH, temperature, or enzyme activity^{[31][32]}. The localized drug depots produced by hydrogel-based systems stick to tissues and provide long-lasting, site-specific release. Numerous benefits are provided by these three-dimensional polymer networks, including methods of controlled release via degradation, diffusion, or routes sensitive to stimuli, Multifunctionality by mixing growth factors or anti-inflammatory drugs with analgesics, Formulations that are injectable and go through sol-to-gel transitions at physiological temperatures^{[34][35][36]}. Nanoparticle Drug Delivery utilizes engineered particles for precise targeting and payload protection. Recent advances include: PLGA nanoparticles with biodegradable properties and BBB penetration capabilities, Functionalized nanoparticles with surface modifications for specific tissue targeting, Combination therapies delivering multiple therapeutic agents simultaneously^{[37][38]}.

Nanoparticle Drug Delivery utilizes engineered particles for precise targeting and payload protection. Recent advances include: PLGA nanoparticles with biodegradable properties and BBB penetration capabilities, Functionalized nanoparticles with surface modifications for specific tissue targeting, Combination therapies delivering multiple therapeutic agents simultaneously. Clinical Applications of targeted delivery systems span multiple routes including perineural injections, intra-articular administration, transdermal patches, and implantable depots, broadening therapeutic applications in both acute and chronic pain management.

These pharmacological advancements collectively represent a paradigm shift toward precision pain management, offering enhanced efficacy, reduced side effects, and improved patient outcomes in pelvic trauma care through mechanism-specific targeting and innovative delivery technologies^[39].

III. REGIONAL ANESTHETIC ADVANCEMENTS

Regional anesthesia has changed pelvic trauma pain management by offering targeted, effective analgesia with fewer systemic effects. Continuous catheter techniques allow long-term administration of local anesthetics, reducing opioid consumption and enabling early mobilization. Techniques such as continuous nerve blocks, especially sacral plexus and pudendal nerve blocks, have demonstrated significant pain reduction and enhanced the functional outcomes [Parker et al., 2010]. The safety and precision of these blocks have been improved by ultrasound guidance and nerve stimulation, allowing for more regular and prolonged analgesia. Depending on the extent of harm and the patterns of fractures, additional regional therapies are also used, such as lumbar plexus blocks and fascia iliaca blocks. In addition to improving the management of pain and lowering hospital stays, minimally invasive surgical techniques in conjunction with regional anesthesia Dabetic & the Associates, 2025. The principal objective of pharmacologic advancements is to enhance analgesia while reducing the hazards related to opioids by using novel medicines and multimodal regimens. With the introduction of imaging and catheter technologies, localized anesthetic procedures have also developed, providing accurate and efficient pain management related to pelvic trauma. The two domains work in tandem to improve outcomes for patients^[40].

1) Nerve Blocks

The use of peripheral and targeted nerve blocks to treat pelvic trauma and persistent pelvic pain has advanced recently. Technique Innovation: With the increased safety and accuracy of ultrasound-guided nerve blocks, medical professionals can target specific pelvic nerves, including the pudendal, obturator, genitofemoral, ilioinguinal, and sacral nerves, for both acute pain relief and diagnostic purposes^[41].

Innovative Strategies: A novel ultrasound-guided treatment for pelvic fracture patients that is based on recent anatomical research is the Pericapsular Nerve Group (PENG) block, which offers targeted pain alleviation^[42].

Clinical Benefits: Especially useful in multimodal pain protocols, nerve block therapies have shown better pain control, fewer opioid need, and fewer systemic side effects than systemic analgesics^[43]. Pudendal and Peripheral Nerve Neuromodulation: Recent research indicates a strong correlation between successful nerve block and pain control with neuromodulation

therapies (miniaturized implantable devices, such as the Freedom® stimulator), opening the door for sustained pain relief where previous interventions have failed. Somatic nerve blocks are used as a diagnostic and therapeutic step for patients with neuropathic or persistent pelvic pain, and they are frequently carried out under CT or ultrasound guidance for maximum selectivity^[44].

2) Epidural Analgesia

Flexible, segmental pain treatment for pelvic injuries is provided by contemporary epidural techniques, because epidural analgesia can be used to titrate sensory blocking and quickly transition to full anesthesia when necessary, it is now used as a primary or adjunctive anesthetic for procedures involving the pelvis, perineum, and lower extremities^[45].

Current technological developments

In contrast to a standard continuous epidural, a dural puncture epidural (DPE) combines the advantages of an epidural with a spinal epidural to give symmetrical, quick pain relief with early sacral coverage and fewer top-ups. Programmed Intermittent Epidural Bolus (PIEB) is compared to continuous infusions, automated intermittent dosing results in better patient satisfaction, less breakthrough pain, less medication intake, and more uniform drug distribution in the epidural region^[46]. The Fig.1.1 clearly shows the techniques of Epidural and Spinal Analgesic.




Fig.1.1: Techniques of neuraxial blockade.

Integration with Multimodal Analgesia: By combining adjuvant medicines (such as opioids, clonidine, and dexmedetomidine) with lower dosages of local anesthetics, epidurals are increasingly used to optimize visceral and somatic pain control while minimizing side effects^[47].

3) Sacral Nerve Blocks

Advanced treatments for pelvic pain are beginning to emerge, including targeted sacral plexus and nerve root blocks, with ultrasound-guided sacral plexus blocks, local anesthetic (with or without steroids) can be precisely applied to the sacral plexus or specific nerve roots (S1–S3), providing notable relief from neuropathic pelvic pain as well as acute injuries^[48]. Sustained decreases in numerical pain scores (NRS) have been demonstrated by repeated ultrasound-guided sacral nerve blocks, these improvements have persisted for months following a course of treatments. These treatments may be especially helpful for people who are not improving with systemic medications or other local methods^[49]. New developments in Sacral neuromodulation, which can be

precisely targeted by imaging and neurophysiological mapping, is an adjuvant for refractory pelvic pain, particularly in disorders like pudendal neuralgia. This broadens its use beyond incontinence^[50].

IV. MINIMALLY INVASIVE AND ADVANCED TECHNIQUES

1) Radiofrequency Ablation (rfa)

After conservative measures fail to alleviate persistent pelvic trauma pain, radiofrequency ablation (RFA) has become a sophisticated minimally invasive technique. The Application and Mechanism in the use of thermal lesioning is by producing heat via radio waves, RFA disrupts the brain's ability to receive pain signals by precisely causing lesions at particular nerve targets^[51]. Peripheral nerve ablation for post-traumatic neuropathic pain syndromes and RFA for lumbar facet joints are examples of recent developments in pelvic applications^[52]. In Innovations technology 63% of patients report pain reductions of at least 50% after using chilled radiofrequency ablation (CRFA), which offers larger, more controlled lesions with long-lasting pain relief for up to 24 months^[52].

2) Spinal Cord Stimulation (SCS)

For refractory pelvic trauma pain, spinal cord stimulation (SCS) is a cutting-edge neuromodulation technique. The Advancements in High-Frequency of SCS at the 12-month follow-up of Ten-kHz technology, 77% of patients in recent prospective multicenter trials report >50% pain reduction, with a 72% decrease in total VAS scores^[53]. Treatment without paraesthesia is the High-frequency stimulation produces analgesic benefits without causing discomfort, although it takes 12 to 48 hours for them to start^[54]. Lead placement improvements: Research indicates that the T8-T9 vertebrae are the best location for bilateral coverage, and in some situations, pain can be reduced by 80% in just two weeks^[54]. **Fig.1.2** shows the spinal cord stimulator system.




Fig.1.2: Spinal cord Stimulator System.

3) Percutaneous Interventions

Advances in Percutaneous Fixation are Decreased morbidity Research shows that, in comparison to non-operative treatment, percutaneous pelvic ring

fixation decreases discomfort, permits faster mobilization, improves the chance of discharge to home, and lowers the risk of one-year mortality^[55]. Technological advancements When compared to traditional methods, modified techniques that use a percutaneous anterior pelvic bridge with K-wire exhibit lower pain scores^[56]. The advantages of an elderly population, Elderly people with fragility fractures benefit most from percutaneous fixation, which speeds up rehabilitation and preserves independence^[57].

V. MULTIMODAL PAIN MANAGEMENT STRATEGIES

1) Combining Modalities

Systematic reviews conducted between 2020 and 2025 show that multimodal analgesia (MMA) significantly lowers postoperative pain scores (WMD = -1.08, $P < 0.001$), shortens hospital stays by 1.19 days, and reduces adverse events by 37%. These findings are supported by meta-analyses^[59]. The MAST Protocol, which combines acetaminophen, naproxen, gabapentin, and lidocaine patches, successfully reduces opioid usage from 48 to 34 MME/day while maintaining comparable pain control in the Multi-modal Analgesic Strategies in Trauma protocol^[21]. The combination techniques synergistic effects target many pain pathways at once, decreasing adverse effects and allowing for lower individual drug dosages^[59].

The two advanced modalities of integration are the physical + psychological and the regional + systemic. When combined with oral/IV analgesics, nerve blocks offer better pain management in the regional + systemic integration than when used alone^[60]. Physical + Psychological When non-invasive non-pharmacological treatments (NINPT) are combined with psychological therapy, the results are improved^[59].

2) Early Mobilization and Rehabilitation

The AO Principles advise 6–12 weeks of non-weight bearing followed by 25% weekly increments in weight load. Evidence-Based Timing in Progressive Weight-Bearing^[61]. The best recovery from pelvic injuries is largely dependent on early mobilization techniques. Following surgery, the protocols permit passive movement in a chair and bed after 15 days, and vigorous rehabilitation can begin 45 days later^[61]. Achieving wheelchair mobility and partial independence through transfers within two to six weeks are short-term objectives during rehabilitation phases^[61]. In the case studies of Physiotherapy Innovations, the multidisciplinary approach shows effective care by early mobilization with an emphasis on range-of-motion exercises, transfer training, and contracture avoidance^[62]. When available, aquatic therapy's integration into water-based rehabilitation offers low-impact, safe exercise^[61]. Advanced rehabilitation regimens for individuals with hip and pelvic fractures provide better results in technology-assisted rehabilitation^[63].

The prevention of complications, which lowers the risk of deep vein thrombosis, pulmonary embolism, skin ulcers, and joint contractures, is one of the clinical benefits of early mobilization and rehabilitation. The functional outcomes include better pain management, an earlier discharge to the patient's home, and lasting functional independence^[64].

3) *Psychological Interventions*

With substantial evidence demonstrating their efficacy, psychological therapies have become essential elements of comprehensive pelvic trauma pain care between 2020 and 2025. Meta-analyses have demonstrated that cognitive behavioral therapy (CBT) improves mental health outcomes while considerably lowering pain levels, including dyspareunia (SMD: -0.54) and dyschezia (MD: -2.90)^[65]. For persistent pelvic pain, Acceptance and Commitment Therapy (ACT), which is especially popular among Australian practitioners, has shown promising results^[66]. Even in cases when symptoms remain, mindfulness-based therapies have been shown to improve quality of life and lessen the severity of pain^[67]. While group therapy programs offer peer support and normalize patient experiences, clinical applications are increasingly embracing trauma-informed care, acknowledging the connection between past physical or emotional trauma and chronic pelvic discomfort^[66]. Single-modality treatments are consistently outperformed by integrated, multidisciplinary strategies that combine medical and psychological therapy^[65].

VI. ADVANCES IN PELVIC TRAUMA PAIN MANAGEMENT

A trend toward multidisciplinary treatments that incorporate pharmacologic, interventional, and rehabilitative strategies has been highlighted by recent studies. With opioids, non-opioid analgesics, and adjuvant drugs based on patient response and the extent of the injury, pharmaceutical treatment is still fundamental. Multimodal pain management, is gaining popularity as a way to lessen opiate dependency and enhance results, Ahmadi and colleagues 2016^[9]. Interventions techniques that target the sacral plexus or pudendal nerve, including nerve blocks, have shown favorable results. For instance, it has been demonstrated that physical therapy and continuous sacral nerve blocks considerably lower pain scores when compared to conservative treatment alone Parker et al., 2010. Based on current trauma care reviews, surgical methods that are minimally invasive for pelvic fractures also result in reduced postoperative pain and a quicker recovery. Dabetic and associates, 2025^{[68][69]}. Rapid evaluation and stabilization within the first 24 hours are essential elements of early care protocols, which aim to minimize secondary injury and improve pain management. The use of external fixation and pelvic binders in damage control orthopedics is essential for minimizing bleeding and the

pain that goes along with it. Incagnoli & Associates, 2019^[70].

VII. CHALLENGES AND BARRIERS

Challenges

The medical picture of pelvic trauma can be extremely complicated, especially when patients are hemodynamically unstable as a result of serious bleeding from damaged pelvic vessels like the venous plexus or internal iliac arteries. Angioembolization or pelvic packing are examples of immediate hemostatic procedures that are commonly necessary, but their prompt implementation is usually impeded by patient instability, diagnostic delays, or restricted access to radiation therapy services in emergency situations. When pelvic trauma happens in conjunction with polytrauma, the situation becomes more complicated since triage and management become more difficult due to concurrent injuries to the chest, abdomen, brain, or spine^[71]. When pelvic trauma happens in conjunction with polytrauma, this complicates matters further because triage and management become more difficult due to the simultaneous injuries to the chest, abdomen, brain, or spine^[72]. Since trauma-related pain can frequently diffuse, referred, or obscured by sedation, shock, or altered state of consciousness, measuring pain is another difficult task that delays the implementation of focused therapies^[73].

Although innovative pain control methods—such as ultrasound-guided nerve blocks, sacral neuromodulation, and transcutaneous electrical nerve stimulation (TENS) have shown promise, their consistent use in acute trauma care remains limited due to a lack of standardized clinical protocols, anatomical variability, and insufficient training among providers Kobayashi et al., 2014^[74]. These challenges are magnified in low-resource settings, where access to imaging technology, surgical tools, and specialized trauma teams is severely restricted Rajasekaran et al., 2020^[74]. Multidisciplinary collaboration is essential but often undermined by role ambiguity, poor communication, and unclear referral pathways between emergency, orthopedic, anesthesia, and rehabilitation teams Falzon et al., 2021^[75]. Moreover, clinicians must strike a balance between adequate pain relief and the risk of masking symptoms of life-threatening complications like compartment syndrome or neurovascular injury^[74]. Despite technological and pharmacological advances, the lack of predictive tools and individualized pain management strategies continues to pose a major challenge in delivering optimal care to pelvic trauma patients^[76].

Pelvic trauma management presents some serious challenges, particularly for patients who are hemodynamically unstable. Since pelvic fractures may cause considerable bleeding, which can harm vital circulatory structures, prompt and precise hemostatic therapies are necessary to prevent evacuation. Additionally, these patients are frequently confused by

several traumas, such as injuries to their chests, abdomens, or brains, which makes it even more challenging to develop treatment plans and do diagnostic examinations. An in-depth analysis by Rajasekaran et al. (Perumal R, S DCR, P SS, Jayaramaraju D, Sen RK, Trikha V) highlights this degree of intricacy. In economically and resource-constrained regions, the proper treatment of pelvic fractures is severely limited by a shortage of advanced medical equipment and a shortage of medical experts with the necessary expertise^[77]. Multidisciplinary Groups Collaboration: For management to be successful, a wide range of healthcare experts must work together in concert. A failure to comprehend roles and referral processes might make collaborative care more difficult. Pelvic trauma presents a highly complex clinical scenario, especially when patients are hemodynamically unstable. One of the primary challenges lies in managing life-threatening hemorrhage due to the disruption of major vessels in the pelvic region, particularly from the venous plexus and internal iliac arteries. Timely bleeding control using interventions like angioembolization or pelvic packing is critical but often delayed due to diagnostic complexity or infrastructural limitations. Moreover, pelvic trauma rarely occurs in isolation patients often present with multiple associated injuries such as abdominal organ damage, thoracic trauma, spinal fractures, or traumatic brain injuries, which significantly complicate triage and treatment decisions. Another challenge is the assessment and management of pain itself. In trauma patients, pain is often diffuse and difficult to localize due to referred pain, altered consciousness, or use of sedatives, making targeted analgesic interventions harder to deliver effectively^[78]. The challenges in managing the pelvic trauma will be explained in **Fig.1.3**.



Fig.1.3: Challenges in effective pelvic trauma care.

Barriers Identified

Numerous significant obstacles prevent the widespread use of novel pain management techniques for pelvic injuries, including neuromodulation, ultrasound-guided nerve blocks, and interdisciplinary care. Since clinicians must also navigate patient-specific anatomical diversity and develop specialized procedural skills which

are frequently lacking due to a lack of training opportunities the lack of standardized clinical protocols for advanced interventions can lead to inconsistent implementation and variable outcomes. The lack of access to sophisticated imaging, specialized equipment, and trained staff further limits prompt intervention and the best possible care for severe pelvic injuries in many healthcare settings, particularly those with minimal resources^[79]. Coordinated pain management between emergency, orthopedic, anesthesia, and rehabilitation teams is often hampered by poor communication, ambiguous team roles, and disjointed referral pathways, which results in treatment that is either delayed or subpar. Effective multidisciplinary collaboration is still a major challenge. These coordination issues are made worse by the dearth of specialized interdisciplinary teams with training in acute pain care and the lack of institutional backing for methodical techniques. Clinicians also constantly struggle to strike a balance between providing strong analgesia and avoiding concealing important signs of complications, like compartment syndrome or neurovascular injury. This leads to cautious prescribing and occasionally insufficient pain management, especially in light of the opioid crisis and medicolegal concerns^[80]. Lastly, it is still challenging to provide patient-centered, efficient management in pelvic trauma care due to the lack of evidence-based guidelines specifically designed for pelvic trauma pain management and healthcare system obstacles like cost concerns, liability issues, and the restricted availability of cutting-edge technologies in settings with limited resources^[81].

Fig.1.4: Factors hindering effective pelvic trauma pain management. **Fig.1.4** will describe the barriers faced in the management of pelvic trauma.




Fig.1.4: Barriers of Pelvic Trauma Management.

VIII. FUTURE DIRECTIONS

Bleeding Control Pre-peritoneal pelvic packing is emphasized as a first-line treatment over angiograms in new policies to achieve quick hemorrhaging control. With this approach, the requirement for big transfusions is reduced and the mortality rate from acute blood loss is decreased ^[82]. Handling Fragility Fractures: Compared to traditional non-surgical approaches, surgical intervention is becoming more and more common for the treatment of fractures of fragility of the pelvis (FFPs), particularly in

older patients. Still, ongoing debates aim to determine the best surgical techniques and patient selection standards^[83]. Standardization of Care is the notable differences in the management of pelvic and acetabular injuries (PAFs) among medical facilities highlight the necessity for consistent treatment procedures. Putting in place a PAF registry could guarantee that therapies are administered consistently and promote treatment improvements^[84]. The use of automated orthopaedic and trauma surgery, including 3D-navigated sacroiliac placement of implants, is improving surgical accuracy and results when treating pelvic fractures^[85]. Robotic surgery is becoming more and more common in female pelvic treatment and reconstructive surgery due to its improved visibility and stitching capabilities in complex pelvic procedures^[86].

IX. CONCLUSION

With the potential for serious bleeding and multi-organ involvement, pelvic trauma is a complex, high-risk disease that requires immediate, multidisciplinary attention. Multidisciplinary pharmacologic regimens, regional anesthetic, and minimally invasive procedures are examples of innovative approaches to pain management which are improving patient outcomes by decreasing opioid dependence and speeding recovery. The broad execution of these innovations is hampered by issues such as inconsistent treatment regimens, high costs, and limited resources. Future study should concentrate on early bleeding management, treatment uniformity, ongoing studies into the effects over time, and the incorporation of cutting-edge methods into all-encompassing, multimodal care networks in order to improve results and guarantee equitable care.

REFERENCES

- [1] Alton, T. B., & Gee, A. O. (2014). Classifications in brief: young and burgess classification of pelvic ring injuries. *Clinical Orthopaedics and Related Research*, 472(8), 2338–2342. <https://doi.org/10.1007/s11999-014-3693-8>.
- [2] Cothren, C. C., Osborn, P. M., Moore, E. E., Morgan, S. J., Johnson, J. L., & Smith, W. R. (2007). Preperitoneal pelvic packing for hemodynamically unstable pelvic fractures: a paradigm shift. *The Journal of Trauma*, 62(4), 834–839; discussion 839–42. <https://doi.org/10.1097/TA.0b013e31803c7632>.
- [3] Madsen, J. E., Flugsrud, G. B., Hammer, N., & Puchwein, P. (2024). Emergency treatment of pelvic ring injuries: state of the art. *Archives of Orthopaedic and Trauma Surgery*, 144(10), 4525–4539. <https://doi.org/10.1007/s00402-024-05447-7>.
- [4] (N.d.).Govfrontiersin.org. Retrieved July 4, 2025, from <http://pubmed.ncbi.nlm.nih.gov/frontiersin.org>.
- [5] (N.d.-b). Nih.gov. Retrieved July 4, 2025, from <http://pmc.ncbi.nlm.nih.gov/>
- [6] PubMed. (n.d.). PubMed. Retrieved July 4, 2025, from <http://pubmed.ncbi.nlm.nih.gov>.
- [7] Altirkistani BA, Ashqar AA, Bahathiq DM, Bougis SM, Aljabri AM, Hanafi S. The effectiveness of ketamine versus opioids in patients with acute pain in the emergency department: A systematic review and meta-analysis. *Cureus* [Internet]. 2023;15(3):e36250. Available from: <http://dx.doi.org/10.7759/cureus.36250>.
- [8] Guay J, Kopp S. Peripheral nerve blocks for hip fractures in adults. *Cochrane Database Syst Rev* [Internet]. 2020;11(4):CD001159. Available from: <http://dx.doi.org/10.1002/14651858.CD001159.pub3>.
- [9] Ahmadi A, Bazargan-Hejazi S, Heidari Zadi Z, Euasobhon P, Ketumarn P, Karbasfrushan A, et al. Pain management in trauma: A review study. *J Inj Violence Res* [Internet]. 2016;8(2):89–98. Available from: <http://dx.doi.org/10.5249/jivr.v8i2.707>.
- [10] Maurice-Szamburski A, Quemeneur C, Rozier R, Cuvillon P, Ecoffey C. Intravenously administered nonsteroidal anti-inflammatory drugs in clinical practice: A narrative review. *Pharmacy (Basel)* [Internet]. 2025;13(1). Available from: <http://dx.doi.org/10.3390/pharmacy13010018>
- [11] Southworth SR, Sellers JA. Narrative summary of recently published literature on intravenous ibuprofen. *Clin Ther* [Internet]. 2020;42(7):1210–21. Available from: <http://dx.doi.org/10.1016/j.clinthera.2020.05.004>
- [12] White PF. The changing role of non-opioid analgesic techniques in the management of postoperative pain. *Anesth Analg* [Internet]. 2005;101(5 Suppl):S5–22. Available from: <http://dx.doi.org/10.1213/01.ANE.0000177099.28914.A7>
- [13] Marchand G, Masoud AT, Govindan M, Ware K, King A, Ruther S, et al. Systematic review and meta-analysis of the efficacy of gabapentin in chronic female pelvic pain without another diagnosis. *AJOG Glob Rep* [Internet]. 2022;2(1):100042. Available from: <http://dx.doi.org/10.1016/j.xagr.2021.100042>
- [14] Lewis SC, Bhattacharya S, Wu O, Vincent K, Jack SA, Critchley HOD, et al. Gabapentin for the management of chronic pelvic pain in women (GaPP1): A pilot randomised controlled trial. *PLoS One* [Internet]. 2016;11(4):e0153037. Available from: <http://dx.doi.org/10.1371/journal.pone.0153037>
- [15] Fda.gov. [cited 2025 Aug 9]. Available from: <https://www.fda.gov/news-events/press->

- announcements/fda-approves-novel-non-opioid-treatment-moderate-severe-acute-pain.
- [16] May M. New approaches to opioid-free pain treatment. *Nat Med* [Internet]. 2025 [cited 2025 Aug 9];31(5):1372–3. Available from: <https://www.nature.com/articles/d41591-025-00024-w>
- [17] Pulskamp TG, Johnson LM, Berlau DJ. Novel non-opioid analgesics in pain management. *Pain Manag* [Internet]. 2024;14(12):641–51. Available from: <http://dx.doi.org/10.1080/17581869.2024.2442292>
- [18] Alessi I, Banton KL, J S, C Z-M, Ch P, Rj R, et al. Exploring novel non-opioid pathways and therapeutics for pain modulation. *Mol Pain* [Internet]. 2025;21:17448069251327840. Available from: <http://dx.doi.org/10.1177/17448069251327840>
- [19] McGovern Medical School. Acute trauma pain multimodal therapy [Internet]. *Surgery*. 2017 [cited 2025 Aug 9]. Available from: <https://med.uth.edu/surgery/acute-trauma-pain-multimodal-therapy/>
- [20] Harvin JA, Albarado R, Truong VTT, Green C, Tyson JE, Pedroza C, et al. Multi-modal analgesic strategy for trauma: A pragmatic randomized clinical trial. *J Am Coll Surg* [Internet]. 2021;232(3):241–251.e3. Available from: <http://dx.doi.org/10.1016/j.jamcollsurg.2020.12.014>
- [21] Sciencedirect.com. [cited 2025 Aug 9]. Available from: <https://www.sciencedirect.com/science/article/abs/pii/S107275152032576X>
- [22] Multimodal pain management: A better approach to pain control - Mayo Clinic [Internet]. *Mayoclinic.org*. [cited 2025 Aug 9]. Available from: <https://www.mayoclinic.org/medical-professionals/trauma/news/multimodal-pain-management-a-better-approach-to-pain-control/mac-20512738>
- [23] Sullivan D, Lyons M, Montgomery R, Quinlan-Colwell A. Exploring opioid-sparing multimodal analgesia options in trauma: A nursing perspective. *J Trauma Nurs* [Internet]. 2016;23(6):361–75. Available from: <http://dx.doi.org/10.1097/JTN.0000000000000250>
- [24] Jamanetwork.com. [cited 2025 Aug 9]. Available from: <https://jamanetwork.com/journals/jama/fullarticle/2796947>
- [25] Gambadoro G, Kopp BJ, Erstad BL. Implications of opioid-sparing medications in critically ill patients: A scoping review. *Med Intensiva (Engl Ed)* [Internet]. 2024;48(12):693–703. Available from: <http://dx.doi.org/10.1016/j.medine.2024.06.020>
- [26] Bhatia A, Buvanendran A. Anesthesia and postoperative pain control-multimodal anesthesia protocol. *J Spine Surg* [Internet]. 2019 [cited 2025 Aug 9];5(Suppl 2):S160–5. Available from: <https://jss.amegroups.org/article/view/4671/html>
- [27] Britishpainsociety.org. [cited 2025 Aug 9]. Available from: https://www.britishpainsociety.org/static/uploads/resources/files/itdd_2015_pro_v3.pdf
- [28] Nhs.uk. [cited 2025 Aug 9]. Available from: <https://www.southtees.nhs.uk/services/pain-management/neuromodulation/intrathecal-drug-delivery-systems/>
- [29] Jain S, Malinowski M, Chopra P, Varshney V, Deer TR. Intrathecal drug delivery for pain management: recent advances and future developments. *Expert Opin Drug Deliv* [Internet]. 2019;16(8):815–22. Available from: <http://dx.doi.org/10.1080/17425247.2019.1642870>
- [30] Nih.gov. [cited 2025 Aug 9]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK538237/>
- [31] Hua S, Wu SY. The use of lipid-based nanocarriers for targeted pain therapies. *Front Pharmacol* [Internet]. 2013;4:143. Available from: <http://dx.doi.org/10.3389/fphar.2013.00143>
- [32] Martinez J, Ingram N, Kapur N, Jayne DG, Beales PA. Vesicle-based formulations for pain treatment: a narrative review. *Pain Rep* [Internet]. 2024;9(6):e1196. Available from: <http://dx.doi.org/10.1097/PR9.00000000000001196>
- [33] Viscusi ER. Liposomal drug delivery for postoperative pain management. *Reg Anesth Pain Med* [Internet]. 2005 [cited 2025 Aug 9];30(5):491–6. Available from: <https://rapm.bmj.com/content/30/5/491>
- [34] Jeong J-O, Kim M, Kim S, Lee KK, Choi H. Advanced hydrogel systems for local anesthetic delivery: Toward prolonged and targeted pain relief. *Gels* [Internet]. 2025;11(2). Available from: <http://dx.doi.org/10.3390/gels11020131>
- [35] Narayanaswamy R, Torchilin VP. Hydrogels and their applications in targeted drug delivery. *Molecules* [Internet]. 2019;24(3):603. Available from: <http://dx.doi.org/10.3390/molecules24030603>
- [36] Li Y, Chen Y, Xue Y, Jin J, Xu Y, Zeng W, et al. Injectable hydrogel delivery system with high drug loading for prolonging local anesthesia. *Adv Sci (Weinh)* [Internet]. 2024;11(24):e2309482. Available from: <http://dx.doi.org/10.1002/advs.202309482>
- [37] Mohammed FS, Omay SB, Sheth KN, Zhou J. Nanoparticle-based drug delivery for the treatment of traumatic brain injury. *Expert Opin Drug Deliv* [Internet]. 2023;20(1):55–73. Available from:

- <http://dx.doi.org/10.1080/17425247.2023.2152001>
- [38] Qin W, Huang J, Zhang M, Xu M, He J, Liu Q. Nanotechnology-based drug delivery systems for treating acute kidney injury. *ACS Biomater Sci Eng* [Internet]. 2024;10(10):6078–96. Available from: <http://dx.doi.org/10.1021/acsbiomaterials.4c01385>
- [39] Raeisi A, Farjadian F. Commercial hydrogel product for drug delivery based on route of administration. *Front Chem* [Internet]. 2024;12:1336717. Available from: <http://dx.doi.org/10.3389/fchem.2024.1336717>
- [40] Mdpi.com. [cited 2025 Jul 10]. Available from: <https://www.mdpi.com/2077-0383/14/5/1492>.
- [41] NYSORA. Ultrasound-guided blocks for pelvic pain [Internet]. NYSORA. 2022 [cited 2025 Aug 9]. Available from: <https://www.nysora.com/pain-management/ultrasound-guided-blocks-for-pelvic-pain/>
- [42] Luftig J, Dreyfuss A, Mantuani D, Howell K, White A, Nagdev A. A new frontier in pelvic fracture pain control in the ED: Successful use of the pericapsular nerve group (PENG) block. *Am J Emerg Med* [Internet]. 2020;38(12):2761.e5-2761.e9. Available from: <http://dx.doi.org/10.1016/j.ajem.2020.05.085>
- [43] Rigaud J, Riant T, Delavierre D, Sibert L, Labat J-J. Les infiltrations des nerfs somatiques dans la prise en charge thérapeutique des douleurs pelvipérinéales chroniques. *Prog Urol* [Internet]. 2010;20(12):1072–83. Available from: <http://dx.doi.org/10.1016/j.purol.2010.08.053>
- [44] Kovacevic N, Sirls L, Gilleran J, Peters K. Peripheral nerve stimulation for pudendal neuralgia and other pelvic pain disorders: current advances. *Front Urol* [Internet]. 2023;3:1323444. Available from: <http://dx.doi.org/10.3389/fruro.2023.1323444>
- [45] Operater. Epidural anesthesia and analgesia [Internet]. NYSORA. 2018 [cited 2025 Aug 9]. Available from: <https://www.nysora.com/topics/abdomen/epidural-anesthesia-analgesia/>
- [46] Analgesia RA in. Novel approaches in regional anesthesia & pain management [Internet]. Openaccessebooks.com. [cited 2025 Aug 9]. Available from: <https://openaccessebooks.com/regional-anesthesia-pain-management/recent-advances-in-labour-analgesia.pdf>
- [47] Bauer M, George JE 3rd, Seif J, Farag E. Recent advances in epidural analgesia. *Anesthesiol Res Pract* [Internet]. 2012;2012:309219. Available from: <http://dx.doi.org/10.1155/2012/309219>
- [48] Ren X-J, Su C-H, Li W, Wang Y. Ultrasound-guided sacral plexus and sacral nerve root blocks alleviating neuropathic pain caused by sacral plexus injury: a case presentation. *Quant Imaging Med Surg* [Internet]. 2024;14(4):3227–30. Available from: <http://dx.doi.org/10.21037/qims-23-1560>
- [49] Domingo-Rufes T, Bong DA, Mayoral V, Ortega-Romero A, Miguel-Pérez M, Sabaté A. Ultrasound-guided pain interventions in the pelvis and the sacral spine. *Tech Reg Anesth Pain Manag* [Internet]. 2013;17(3):107–30. Available from: <http://dx.doi.org/10.1053/j.trap.2014.01.014>
- [50] Perrouin-Verbe M-A, Van Kerrebroeck PEV. The future of neuromodulation for functional pelvic problems. *Continence (Amst)* [Internet]. 2024;11(101694):101694. Available from: <http://dx.doi.org/10.1016/j.cont.2024.101694>
- [51] Radiofrequency ablation (RFA) is a procedure that uses heat to destroy tissue. For pain management involving joints, RFA heats a nerve, which stops or reduces pain signals to your brain [Internet]. Cleveland Clinic. 2017 [cited 2025 Aug 9]. Available from: <https://my.clevelandclinic.org/health/treatments/17411-radiofrequency-ablation>
- [52] Abd-Elsayed A, Robinson CL, Peters T. Narrative review of radiofrequency ablation applications in peripheral nerves. *Ann Palliat Med* [Internet]. 2024 [cited 2025 Aug 9];13(4):893–900. Available from: <https://apm.amegroups.org/article/view/124175/html>
- [53] Patel CB, Patel AA, Diwan S. Pain Physician 2022; 25:E531-E542 • ISSN 2150-1149 [Internet]. Painphysicianjournal.com. [cited 2025 Aug 9]. Available from: <https://www.painphysicianjournal.com/current/pdf?article=NzQ4MQ%3D%3D&journal=144>
- [54] Romero-Serrano E, Esparza-Miñana JM. Spinal cord stimulation in the approach to chronic pelvic pain: A case report and literature review: A case report and literature review. *Medicine (Baltimore)* [Internet]. 2021;100(52):e28379. Available from: <http://dx.doi.org/10.1097/MD.00000000000028379>
- [55] de Ridder VA, Whiting PS, Balogh ZJ, Mir HR, Schultz BJ, Routt MC. Pelvic ring injuries: recent advances in diagnosis and treatment. *OTA Int* [Internet]. 2023;6(3 Suppl):e261. Available from: <http://dx.doi.org/10.1097/OI9.0000000000000261>
- [56] Huang G, Cai L, Jia X, Ji Y, Zhou Y, Mou X, et al. A novel hybrid fixation (percutaneous anterior pelvic bridge with K-wire) for the treatment of traumatic pelvic ring injury: A prospective study. *Int J Surg* [Internet]. 2019;61:11–6. Available from: <http://dx.doi.org/10.1016/j.ijsu.2018.11.025>
- [57] Clinic C. Percutaneous fixation: An underused treatment for pelvic ring fractures in older adults [Internet]. Cleveland Clinic. 2025 [cited 2025 Aug 9]. Available from:

- <https://consultqd.clevelandclinic.org/percutaneous-pelvic-ring-fixation-in-older-adults>
- [58] Nature.com. [cited 2025 Aug 9]. Available from: <https://www.nature.com/articles/s41598-025-00069-4>
- [59] Shi Y, Wu W. Multimodal non-invasive non-pharmacological therapies for chronic pain: mechanisms and progress. *BMC Med* [Internet]. 2023;21(1):372. Available from: <http://dx.doi.org/10.1186/s12916-023-03076-2>
- [60] Sampognaro G, Harrell R. Multimodal postoperative pain control after orthopaedic surgery. In: *StatPearls*. Treasure Island (FL): StatPearls Publishing; 2025.
- [61] Piccione F, Maccarone MC, Cortese AM, Rocca G, Sansubrinio U, Piran G, et al. Rehabilitative management of pelvic fractures: a literature-based update. *Eur J Transl Myol* [Internet]. 2021;31(3). Available from: <http://dx.doi.org/10.4081/ejtm.2021.9933>
- [62] Patil DS, Thakre VM, Gangwani N. Physiotherapy management focusing on early mobilisation in complex pelvic and pott's fracture. *Cureus* [Internet]. 2023;15(11):e49525. Available from: <http://dx.doi.org/10.7759/cureus.49525>
- [63] Katz D, Gefitler A, Abu-Ajaj A, Makulin E, Star E, Zikrin E, et al. Intensive rehabilitation after pelvic and hip fractures: a comparative retrospective study. *Front Med (Lausanne)* [Internet]. 2024;11:1346354. Available from: <http://dx.doi.org/10.3389/fmed.2024.1346354>
- [64] Mukhopadhyay R. Role of rehabilitation after pelvic and acetabular surgery [Internet]. Dr. Reet Mukhopadhyay -. Dr. Reet Mukhopadhyay; 2024 [cited 2025 Aug 9]. Available from: <https://orthodocreet.com/role-of-rehabilitation-after-pelvic-and-acetabular-surgery/>
- [65] Brooks T, Sharp R, Evans S, Baranoff J, Esterman A. Psychological interventions for women with persistent pelvic pain: A survey of mental health clinicians. *J Multidiscip Healthc* [Internet]. 2021;14:1725–40. Available from: <http://dx.doi.org/10.2147/JMDH.S313109>
- [66] Clinic C. Treatment for the psychological component of chronic pelvic pain [Internet]. Cleveland Clinic. 2025 [cited 2025 Aug 9]. Available from: <https://consultqd.clevelandclinic.org/the-psychological-component-of-chronic-pelvic-pain>
- [67] Del Pino-Sedeño T, Cabrera-Maroto M, Abrante-Luis A, González-Hernández Y, Ortíz Herrera MC. Effectiveness of psychological interventions in endometriosis: a systematic review with meta-analysis. *Front Psychol* [Internet]. 2024;15:1457842. Available from: <http://dx.doi.org/10.3389/fpsyg.2024.1457842>
- [68] Ireland E, Womack J, Foley E. Sacral plexus catheter analgesia in a patient with traumatic sacral fractures and Tarlov cysts. *Anaesth Rep* [Internet]. 2019;7(2):76–8. Available from: <http://dx.doi.org/10.1002/anr3.12023>
- [69] Mdpi.com. [cited 2025 Jul 10]. Available from: <https://www.mdpi.com/2077-0383/14/5/1492>.
- [70] Incagnoli P, Puidupin A, Ausset S, Beregi JP, Bessereau J, Bobbia X, et al. Early management of severe pelvic injury (first 24 hours). *Anaesth Crit Care Pain Med* [Internet]. 2019;38(2):199–207. Available from: <http://dx.doi.org/10.1016/j.accpm.2018.12.003>.
- [71] Osborn PM, Sathy AK, Starnes BW. Hemorrhage control in pelvic fractures. *Semin Intervent Radiol*. 2020;37(1):48–56
- [72] Madhu TS, Raman R, Giannoudis PV. Long-term outcome in patients with combined spinal and pelvic fractures. *Injury* [Internet]. 2007;38(5):598–606. Available from: <http://dx.doi.org/10.1016/j.injury.2006.11.005>
- [73] Cohen SP, Christo PJ, Moroz L. Pain management in trauma patients. *Am J Phys Med Rehabil* [Internet]. 2004 [cited 2025 Jul 16];83(2):142–61. Available from: https://journals.lww.com/ajpmr/abstract/2004/02000/pain_management_in_trauma_patients.11.aspx
- [74] Fabbri A, Voza A, Riccardi A, Serra S, Iaco FD, Study and Research Center of the Italian Society of Emergency Medicine (SIMEU). The pain management of trauma patients in the emergency department. *J Clin Med* [Internet]. 2023;12(9). Available from: <http://dx.doi.org/10.3390/jcm12093289>
- [75] Ding H, Wang W, Yuan M, Tang J, Zhou J, Pan J. Multidisciplinary management of severe open pelvic fracture with multiple organ injuries: A case report. *Medicine (Baltimore)* [Internet]. 2025;104(31):e43551. Available from: <http://dx.doi.org/10.1097/MD.00000000000043551>
- [76] Siqueira-Campos VM, de Deus MSC, Poli-Neto OB, Rosa-E-Silva JC, de Deus JM, Conde DM. Current challenges in the management of chronic pelvic pain in women: From bench to bedside. *Int J Womens Health* [Internet]. 2022;14:225–44. Available from: <http://dx.doi.org/10.2147/IJWH.S224891>
- [77] Researchgate.net. [cited 2025 Jul 10]. Available from: https://www.researchgate.net/publication/365129283_The_management_of_pelvic_ring_fractures_in_low-resource_environments_review.
- [78] Sobantu NA, Tshabalala MD, Chetty V. Exploring the collaborative care of patients with pelvic fractures in Tshwane, South Africa. *S Afr Fam Pract* (2004) [Internet]. 2023;65(1):e1–9. Available from: <http://dx.doi.org/10.4102/safp.v65i1.5705>.
- [79] Nagpal AK, Gadkari C, Singh A, Pundkar A. Optimizing pain management in Emergency

- Departments: A comprehensive review of current analgesic practices. Cureus [Internet]. 2024;16(9):e69789. Available from: https://assets.cureus.com/uploads/review_article/pdf/299073/20241020-1706737-17tn37.pdf
- [80] Nożewski J, Bondarczuk R, Hołody G, Kitt M, Mućka N, Religioni U. Barriers to improving pain management in the emergency department: Lessons from a lean-driven quality improvement initiative. *J Clin Med* [Internet]. 2025;14(13). Available from: <http://dx.doi.org/10.3390/jcm14134566>
- [81] Nyirigira G, Rutayisire F, Ndekezi JK, Kwizera R, Neil KL. Barriers and facilitators to implementing acute pain services at King Faisal Hospital Rwanda. *Pain Res Manag* [Internet]. 2025;2025:8360891. Available from: <http://dx.doi.org/10.1155/prm/8360891>
- [82] Parry JA, Smith WR, Moore EE, Burlew CCC, Mauffrey C. The past, present, and future management of hemodynamic instability in patients with unstable pelvic ring injuries. *Injury* [Internet]. 2021;52(10):2693–6. Available from: <http://dx.doi.org/10.1016/j.injury.2020.02.101>
- [83] Hutchings L, Roffey DM, Lefaivre KA. Fragility fractures of the pelvis: Current practices and future directions. *Curr Osteoporos Rep* [Internet]. 2022;20(6):469–77. Available from: <http://dx.doi.org/10.1007/s11914-022-00760-9>
- [84] Martin A, McMaster J, Bretherton C, Noyes D. Pelvic and acetabular fracture care in England: current workload and future directions. *Ann R Coll Surg Engl* [Internet]. 2021;103(6):420–5. Available from: <http://dx.doi.org/10.1308/rcsann.2021.0015>
- [85] Stübig T, Windhagen H, Krettek C, Ettinger M. Computer-assisted orthopedic and trauma surgery. *Dtsch Arztebl Int* [Internet]. 2020;117(47):793–800. Available from: <http://dx.doi.org/10.3238/arztebl.2020.0793>
- [86] Urologyresearchandpractice.org. [cited 2025 Jul 10]. Available from: <https://www.urologyresearchandpractice.org/en/the-current-state-and-the-future-of-robotic-surgery-in-female-pelvic-medicine-and-reconstructive-surgery-133520>