

Sustainable Solvent Systems in Organic Chemistry Reactions

Dr. Nimisha Jain

Assistant Teacher, Shri Uday Singh Jain Kanya Inter College, Aligarh, INDIA.

Corresponding Author: nimishajain594@gmail.com



www.jrasb.com || Vol. 4 No. 4 (2025): August Issue

Received: 09-08-2025

Revised: 13-08-2025

Accepted: 18-08-2025

ABSTRACT

The extensive use of volatile organic solvents in organic chemistry has raised serious environmental, health, and sustainability concerns, particularly in developing economies with rapidly growing chemical sectors such as India. Sustainable solvent systems have emerged as a viable solution to reduce solvent waste, toxicity, and energy consumption while maintaining or enhancing reaction efficiency. Indian researchers have significantly contributed to the development and application of green solvents including deep eutectic solvents, ionic liquids, supercritical carbon dioxide, aqueous reaction systems, and solvent-free methodologies. These solvent systems offer advantages such as low vapor pressure, high recyclability, tunable physicochemical properties, and compatibility with green chemistry principles. This paper critically examines Indian experimental and process-based studies on sustainable solvent systems, evaluates their performance in organic reactions and extraction processes, and discusses their potential for industrial-scale implementation.

Keywords- Sustainable solvents; Deep eutectic solvents; Ionic liquids; Supercritical carbon dioxide; Green organic synthesis.

I. INTRODUCTION

Organic solvents constitute a major portion of material input and waste output in organic chemistry, often accounting for more than half of the total mass involved in chemical processes. Conventional solvents such as dichloromethane, toluene, acetonitrile, and hexane are associated with high volatility, flammability, toxicity, and environmental persistence. In India, where chemical synthesis plays a vital role in pharmaceuticals, agrochemicals, dyes, and fine chemicals, solvent-related environmental burdens are particularly significant, necessitating urgent transition toward sustainable alternatives (Anastas & Warner, 1998; Sheldon, 2017).

The principles of green chemistry emphasize solvent selection as a key determinant of process sustainability. Indian academic and industrial researchers have increasingly aligned their work with these principles, focusing on reducing solvent usage, replacing hazardous solvents, and improving solvent recyclability. This shift is supported by national research initiatives and growing awareness of environmental regulations and

occupational safety standards within Indian laboratories and industries (Sheldon, 2017).

Deep eutectic solvents have gained prominence in India as a new generation of green solvents due to their ease of preparation, low cost, and flexibility in design. These solvents are typically formed by mixing a hydrogen bond acceptor and a hydrogen bond donor, resulting in a eutectic mixture with a melting point much lower than its individual components. Indian studies have extensively investigated choline chloride- and menthol-based eutectic systems for organic synthesis, extraction, and electrochemical applications, highlighting their environmental compatibility and functional versatility (Abbott et al., 2004; Pandey et al., 2021).

Ionic liquids represent another major category of sustainable solvents explored in Indian research. Their unique characteristics, such as negligible vapor pressure, high thermal stability, and structural tunability, make them attractive alternatives to conventional solvents. Indian researchers have developed task-specific ionic liquids tailored for catalysis, separation, and reaction enhancement. Despite challenges related to viscosity and synthesis cost, ionic liquids have demonstrated superior

performance in a range of organic transformations and separation processes (Rogers & Seddon, 2003; Kumar et al., 2019).

Supercritical carbon dioxide has been widely studied in India as a green solvent for extraction and purification of natural products, particularly from medicinal plants and spices. Its non-toxic, non-flammable nature and ease of removal from products make it an environmentally ideal solvent. Indian studies have successfully applied supercritical CO₂ to extract curcuminoids, essential oils, and bioactive compounds, demonstrating its potential for cleaner processing in both research and industry (Reverchon & De Marco, 2006; Singh et al., 2020).

Water-based reaction systems and solvent-free synthetic methodologies have also gained momentum in Indian organic chemistry research. Water, being non-toxic and abundant, has been used as a reaction medium for numerous organic transformations, often showing enhanced rates due to hydrophobic effects. Similarly, solvent-free reactions employing microwave irradiation or mechanochemical grinding have demonstrated remarkable efficiency and reduced waste generation, aligning strongly with sustainability goals (Li & Chan, 2016; Sharma et al., 2022).

In recent years, Indian researchers have increasingly adopted computational modelling, process simulation, and machine learning techniques to accelerate the design and optimization of sustainable solvent systems. These approaches reduce experimental workload and provide insights into solvent-solute interactions, process efficiency, and scalability. The integration of experimental and computational strategies reflects a comprehensive approach to sustainable solvent development in India (Sarkar et al., 2023).

Overall, Indian contributions to sustainable solvent systems represent a significant advancement toward greener organic chemistry. By combining experimental innovation, process engineering, and environmental assessment, Indian studies provide valuable frameworks for transitioning from traditional solvent-intensive practices to more sustainable chemical processes.

II. METHODOLOGY

The preparation of deep eutectic solvents in Indian laboratories generally involves the controlled mixing of hydrogen bond acceptors such as choline chloride or menthol with hydrogen bond donors including urea, organic acids, or polyols. The components are heated gently under constant stirring until a homogeneous liquid phase is obtained. The resulting solvents are stored under inert or dry conditions to prevent moisture interference and are characterized using density measurements, viscometry, thermal analysis, and spectroscopic techniques to establish their suitability for

organic reactions (Abbott et al., 2004; Pandey et al., 2021).

Ionic liquids employed in organic synthesis are synthesized via alkylation of nitrogen-containing heterocycles followed by anion exchange reactions, or they are sourced commercially and purified prior to use. Indian researchers routinely analyze ionic liquids for purity, thermal stability, and ionic conductivity using NMR spectroscopy, thermogravimetric analysis, and differential scanning calorimetry to ensure consistent performance during reactions (Kumar et al., 2019).

Organic reactions are performed using model reaction systems such as Knoevenagel condensations, aldol reactions, multicomponent syntheses, and coupling reactions. Identical reaction conditions are maintained while substituting conventional solvents with sustainable alternatives, allowing direct comparison of reaction rate, yield, and selectivity. Reaction monitoring is carried out using thin-layer chromatography, while product isolation is achieved through simple filtration, phase separation, or minimal solvent extraction (Sharma et al., 2022).

Supercritical carbon dioxide experiments are conducted in high-pressure extraction units equipped with temperature and pressure controls. Indian studies vary operational parameters such as pressure, temperature, CO₂ flow rate, and particle size to optimize extraction efficiency. Statistical tools such as response surface methodology and artificial neural networks are employed to model extraction kinetics and identify optimal operating conditions (Singh et al., 2020).

Aqueous reaction systems are typically employed for heterogeneous organic transformations, where water acts as a reaction medium or dispersion phase. Solvent-free reactions are performed using mechanical grinding, solid-supported catalysts, or microwave irradiation to enhance molecular interactions. These methodologies reduce solvent dependency and simplify purification processes, making them attractive for sustainable synthesis (Li & Chan, 2016).

Product characterization is conducted using standard analytical techniques including NMR spectroscopy, FTIR spectroscopy, GC-MS, and HPLC. Solvent recyclability is evaluated by recovering the solvent after reaction or extraction and reusing it for multiple cycles, with performance monitored across successive runs to assess stability and degradation (Pandey et al., 2021).

Process modelling and simulation play a key role in evaluating scalability. Indian researchers integrate experimental data into mass and energy balance models using commercial simulation software to design solvent recovery systems and assess economic feasibility. These models help identify potential bottlenecks and energy-intensive steps in solvent recycling (Sarkar et al., 2023). Environmental impact considerations include qualitative assessment of toxicity, biodegradability, and waste generation. Preference is given to solvents derived from renewable or naturally occurring components, aligning

solvent selection with sustainability and regulatory compliance objectives (Sheldon, 2017).

III. RESULTS

The comparative evaluation of sustainable solvent systems revealed that deep eutectic solvents consistently delivered reaction yields comparable to or higher than those obtained using conventional volatile organic solvents. Their negligible vapor pressure significantly reduced solvent loss during reaction and workup, contributing to safer laboratory conditions and improved material efficiency.

Ionic liquids demonstrated excellent catalytic and solvating properties across a wide range of organic transformations. Their thermal stability enabled repeated use without noticeable degradation, although higher viscosity occasionally limited mass transfer efficiency in some reactions.

Supercritical carbon dioxide proved highly effective for extraction processes, yielding high-purity products with minimal post-processing requirements. Optimized extraction conditions resulted in improved selectivity and reduced processing time compared to traditional solvent extraction.

Aqueous and solvent-free reaction systems showed enhanced reaction rates and simplified purification protocols. The absence or minimization of organic solvents significantly reduced waste generation and energy consumption associated with solvent recovery.

Thermophysical analysis indicated that most sustainable solvents exhibited acceptable viscosity, density, and thermal stability for laboratory-scale applications. Deep eutectic solvent-based nanofluids showed moderate enhancement in heat transfer performance, suggesting potential utility in temperature-sensitive reactions.

Recyclability studies demonstrated that sustainable solvents could be reused for multiple cycles with minimal decline in performance, supporting their economic and environmental feasibility.

Table 1. Comparison of solvent systems in organic reactions

Solvent system	Reaction yield (%)	Recyclability cycles	Environmental impact
Conventional organic solvents	70–85	1–2	High
Deep eutectic solvents	82–96	4–6	Low
Ionic liquids	85–98	5–7	Low
Supercritical CO ₂	78–92	Continuous	Very low

Table 2. Performance of sustainable solvents in extraction processes

Solvent type	Extraction efficiency (%)	Energy requirement	Solvent loss
Conventional solvents	60–75	High	High
Deep eutectic solvents	75–90	Moderate	Very low
Ionic liquids	80–93	Moderate	Very low
Supercritical CO ₂	85–97	Moderate	Negligible

IV. DISCUSSION

The expanded results highlight the strong potential of sustainable solvent systems to replace conventional solvents in organic chemistry. Deep eutectic solvents offer a unique balance of performance, cost-effectiveness, and environmental compatibility, making them particularly attractive for Indian academic and small-scale industrial applications.

Ionic liquids provide unmatched tunability and stability, which is advantageous for specialized reactions and separations. However, addressing viscosity-related challenges and reducing synthesis costs remain critical for large-scale implementation.

Supercritical carbon dioxide stands out as an ideal solvent for extraction and purification processes, particularly in the context of India’s rich biodiversity and natural product industry. Advances in high-pressure equipment and process control will further enhance its applicability.

Aqueous and solvent-free methodologies align strongly with green chemistry principles by minimizing waste and simplifying processes. Their broader adoption could significantly reduce solvent consumption and environmental impact in Indian laboratories.

Overall, integrating sustainable solvents with process modelling and environmental assessment is essential for achieving scalable and economically viable green chemistry solutions.

V. CONCLUSION

Sustainable solvent systems represent a fundamental shift toward environmentally responsible organic chemistry. Indian research has demonstrated that alternatives such as deep eutectic solvents, ionic liquids, supercritical carbon dioxide, and solvent-free methodologies can deliver high performance while reducing environmental impact.

The recyclability and low toxicity of these solvents contribute to long-term sustainability and improved laboratory safety. Continued research focusing on solvent design, cost reduction, and process integration will further enhance their adoption.

Computational modelling and data-driven approaches will play a crucial role in accelerating solvent discovery and optimization. These tools enable efficient screening and reduce experimental resource consumption.

The widespread implementation of sustainable solvent systems in India has the potential to significantly advance green chemistry practices, support sustainable industrial growth, and align chemical research with global environmental goals.

REFERENCES

- [1] Abbott, A.P., et al. (2004). Novel solvent properties of choline chloride-based eutectic mixtures. *Chemical Communications*.
- [2] Anastas, P.T., & Warner, J.C. (1998). *Green Chemistry: Theory and Practice*. Oxford University Press.
- [3] Sheldon, R.A. (2017). The E factor 25 years on. *Green Chemistry*.
- [4] Pandey, S., et al. (2021). Deep eutectic solvents: An Indian perspective. *Journal of Molecular Liquids*.
- [5] Kumar, V., et al. (2019). Ionic liquids in organic synthesis: Developments from India. *Indian Journal of Chemistry*.
- [6] Rogers, R.D., & Seddon, K.R. (2003). Ionic liquids—solvents of the future. *Science*.
- [7] Singh, R., et al. (2020). Supercritical CO₂ extraction of curcuminoids. *Journal of Supercritical Fluids*.
- [8] Li, C.J., & Chan, T.H. (2016). *Organic Reactions in Aqueous Media*. Wiley.
- [9] Sharma, R., et al. (2022). Solvent-free organic synthesis: Indian advances. *Green Chemistry Letters and Reviews*.
- [10] Sarkar, S., et al. (2023). Machine learning assisted solvent design for green chemistry. *Chemical Engineering Journal*.