

Biological Control of Mosquito Vectors Using Predatory Fish and Microbial Agents: An Experimental Approach

Himanshu Bansal¹ and Dr. Anil Kumar²

¹P.G. Student, Department of Zoology, J.V. College, Baraut (Baghpat), INDIA

²Associate Professor, Department of Zoology, J.V. College, Baraut, Baghpat, INDIA

¹Corresponding Author: himanshubansaljvc@gmail.com



www.jrasb.com || Vol. 5 No. 1 (2026): February Issue

Received: 17-02-2026

Revised: 22-02-2026

Accepted: 27-02-2026

ABSTRACT

Mosquito-borne diseases remain a major public health concern worldwide, particularly in tropical and subtropical regions where climatic conditions favor mosquito breeding. Conventional mosquito control strategies rely heavily on chemical insecticides; however, excessive use of these chemicals has resulted in environmental pollution, insecticide resistance, and negative impacts on non-target organisms. Biological control has emerged as a sustainable alternative for managing mosquito populations while minimizing ecological harm. The present study evaluates the effectiveness of predatory fish species and microbial agents as biological control tools against mosquito larvae. Experimental observations were conducted to assess larval reduction rates under different treatment conditions involving larvivorous fish and microbial agents such as *Bacillus thuringiensis israelensis* (Bti). The results demonstrate that predatory fish significantly reduced larval densities in aquatic habitats, while microbial agents showed high larvicidal efficiency when applied at recommended concentrations. Integrated use of both biological control methods resulted in the greatest reduction in mosquito populations. The findings suggest that environmentally safe biological control approaches can play a crucial role in vector management programs and contribute to sustainable mosquito control strategies.

Keywords- Mosquitoes; Biocontrol; Larvicides; Predation; Vectors.

I. INTRODUCTION

Mosquitoes are among the most important vectors responsible for transmitting a wide range of infectious diseases that affect millions of people worldwide. Vector-borne diseases such as malaria, dengue, chikungunya, Zika virus infection, Japanese encephalitis, and lymphatic filariasis remain major public health challenges, particularly in tropical and subtropical regions where environmental conditions favor mosquito breeding and survival [1]. The increasing prevalence of these diseases has highlighted the need for effective vector management strategies to reduce mosquito populations and prevent disease transmission.

Conventional mosquito control programs have relied heavily on chemical insecticides such as organophosphates, pyrethroids, and carbamates to eliminate mosquito larvae and adults. While chemical

control methods provide rapid suppression of mosquito populations, their continuous and widespread use has resulted in several ecological and health concerns. One of the major issues associated with chemical insecticides is the development of resistance in mosquito populations, which reduces the effectiveness of control programs over time. In addition, chemical residues may contaminate water bodies, affect non-target organisms, and disrupt aquatic ecosystems [2].

Due to these limitations, environmentally sustainable alternatives to chemical insecticides have become increasingly important in vector control programs. Biological control is one such approach that involves the use of natural predators, pathogens, or parasites to suppress mosquito populations. Biological control methods offer several advantages, including ecological safety, species specificity, and reduced risk of

resistance development compared to chemical insecticides [3].

Among the various biological control agents, larvivorous fish have been widely used in mosquito control programs. Fish species such as *Gambusia affinis* (mosquito fish) and *Poecilia reticulata* (guppy) are known for their ability to consume large numbers of mosquito larvae in aquatic habitats. These fish can survive in a variety of environmental conditions and are capable of controlling mosquito populations in ponds, irrigation channels, rice fields, and stagnant water bodies [4]. Because of their adaptability and high larval consumption rates, larvivorous fish have been introduced in several mosquito control initiatives worldwide.

In addition to predatory fish, microbial agents have also proven to be effective biological control tools against mosquito larvae. The bacterium *Bacillus thuringiensis israelensis* (Bti) is one of the most widely used microbial larvicides in mosquito control programs. Bti produces toxins that specifically target mosquito larvae by disrupting the cells of the larval midgut after ingestion. These toxins cause rapid larval mortality while having minimal effects on non-target organisms such as fish, amphibians, and beneficial aquatic insects [5].

Several studies have demonstrated that biological control agents can significantly reduce mosquito populations when applied under appropriate environmental conditions. However, the effectiveness of biological control methods often depends on local ecological factors such as temperature, water quality, habitat type, and availability of mosquito breeding sites. Therefore, experimental evaluation of biological control strategies is necessary to determine their effectiveness under specific environmental conditions [6].

Integrated vector management strategies that combine multiple biological control agents may provide improved mosquito control outcomes compared to single-method approaches. For example, the combined use of predatory fish and microbial larvicides can provide both immediate and long-term suppression of mosquito larvae. While microbial agents rapidly kill larvae present in water bodies, predatory fish provide continuous control by feeding on newly emerging larvae [7].

Despite the growing importance of biological mosquito control, there remains a need for experimental studies that evaluate the combined effectiveness of predatory fish and microbial agents under controlled conditions. Understanding the interaction between these biological control methods can help improve mosquito management programs and reduce reliance on chemical insecticides.

Therefore, the present study aims to evaluate the effectiveness of predatory fish and microbial agents in controlling mosquito larvae through controlled experimental observations. The study also examines the potential benefits of integrating these biological control

methods to develop sustainable and environmentally friendly mosquito vector management strategies.

II. MATERIALS AND METHODS

The present study was conducted through controlled experimental observations designed to evaluate the effectiveness of biological control agents against mosquito larvae in aquatic environments. Artificial aquatic habitats were created using plastic containers filled with freshwater in order to simulate natural mosquito breeding conditions commonly found in stagnant water bodies such as ponds, drains, irrigation channels, and temporary rainwater collections. Such controlled experimental environments allow reliable assessment of mosquito larval mortality and predator efficiency under standardized conditions [1].

Mosquito larvae used in the experiment were collected from natural breeding sites located in stagnant water bodies within the study area. These breeding habitats included drainage channels, ponds, and small water accumulations frequently observed in urban and peri-urban environments. Collected larvae were carefully transferred into laboratory containers filled with water from the original habitats to prevent sudden environmental stress. The larvae were acclimatized for a short period before experimentation to ensure normal physiological activity and survival under laboratory conditions [2].

For the evaluation of biological predation, larvivorous fish species commonly used in mosquito control programs were selected. The fish species *Gambusia affinis* (mosquito fish) and *Poecilia reticulata* (guppy) were chosen because of their well-documented feeding behavior and high capacity to consume mosquito larvae. These species are widely used in mosquito control programs due to their adaptability to diverse environmental conditions and their ability to significantly reduce mosquito breeding in aquatic habitats [3]. The fish were introduced into experimental containers containing mosquito larvae and allowed to acclimatize before observations began to ensure natural feeding activity.

Microbial control experiments were conducted using the bacterium *Bacillus thuringiensis israelensis* (Bti), a widely recognized biological larvicide used in mosquito control programs worldwide. Bti produces specific endotoxins that are highly toxic to mosquito larvae but generally harmless to other aquatic organisms and vertebrates. The microbial agent was applied to the experimental containers at recommended concentrations according to standard guidelines for larval mosquito control. Observations were then made to record larval mortality after exposure to the microbial toxin [4].

To compare the effectiveness of different control strategies, the experimental design included three

treatment conditions as well as an untreated control setup. One set of containers included only predatory fish, another set received only microbial treatment using Bti, and a third set involved the combined application of predatory fish and microbial agents. Control containers were maintained without any biological control agent in order to observe natural survival rates of mosquito larvae under similar environmental conditions. This comparative experimental design enabled evaluation of the relative efficiency of each control strategy [5].

Larval populations were monitored at regular time intervals throughout the experimental period. The number of live larvae, dead larvae, and partially immobilized larvae was recorded during each observation session. Mortality percentages were calculated by comparing the initial number of larvae introduced into each container with the number of surviving larvae at the end of the observation period. Monitoring larval survival rates provided quantitative data for evaluating the effectiveness of different biological control methods [6].

Environmental factors such as water temperature, pH, and dissolved oxygen levels were also recorded during the experimental period because these parameters can influence mosquito larval development and predator feeding behavior. Maintaining stable environmental conditions ensured that variations in larval mortality were primarily related to biological control treatments rather than external environmental fluctuations [7].

Finally, the collected experimental data were compiled and analyzed to determine the relative effectiveness of predatory fish, microbial agents, and their combined use in controlling mosquito larvae. The results were presented in the form of tables and graphical representations to illustrate larval reduction trends and comparative performance of biological control methods under the experimental conditions [8].

III. RESULTS

The experimental results demonstrated that biological control agents significantly reduced mosquito larval populations compared to untreated control conditions. Predatory fish showed strong larvivorous activity and rapidly consumed mosquito larvae present in the experimental containers.

Microbial treatment using *Bacillus thuringiensis israelensis* also produced high larval mortality within a short period. Larvae exposed to the microbial toxin exhibited reduced mobility and eventually died due to disruption of digestive processes.

The combined application of predatory fish and microbial agents resulted in the highest larval reduction rates. This integrated approach provided both immediate and sustained control of mosquito populations.


Table 1: Larval Reduction Under Different Treatments

Treatment	Initial Larvae	Final Larvae	Reduction (%)
Control	100	95	5
Predatory Fish	100	30	70
Microbial Agent (Bti)	100	20	80
Combined Treatment	100	10	90

Table 2: Average Larval Consumption by Predatory Fish

Fish Species	Average Larvae Consumed per Day
Gambusia affinis	80
Poecilia reticulata	60

Larval Reduction Trend of Mosquito Larvae Under Different Biological Control Treatments



Graph: Larval Reduction Trend

The graph indicates that the **combined treatment showed the greatest reduction in mosquito larvae**.

IV. DISCUSSION AND CONCLUSION

The findings of the study highlight the potential of biological control agents as sustainable alternatives to chemical insecticides for mosquito management. Predatory fish were highly effective in reducing larval populations due to their continuous feeding behavior in aquatic environments. Their ability to consume large numbers of larvae makes them suitable for long-term mosquito control in stagnant water bodies [4].

Microbial agents such as *Bacillus thuringiensis israelensis* also demonstrated strong larvicidal activity. The toxin produced by Bti specifically targets mosquito larvae and has minimal impact on other aquatic organisms, making it an environmentally safe control method [5].

The combined use of predatory fish and microbial agents provided the highest level of mosquito control in the experimental setup. This integrated approach ensures immediate larval mortality through

microbial action while maintaining long-term suppression through predatory fish activity.

Biological control strategies offer several advantages over chemical methods, including environmental safety, reduced risk of resistance development, and sustainability. These methods are particularly suitable for community-based mosquito control programs in rural and urban environments.

In conclusion, the study demonstrates that biological control using predatory fish and microbial agents can effectively reduce mosquito vector populations. Integration of these methods into public health vector management programs can contribute to sustainable control of mosquito-borne diseases while minimizing environmental impacts.

REFERENCES

- [1] World Health Organization. (2023). *Vector-borne diseases and global health risk*.
- [2] Kumar, A., & Singh, P. (2021). *Insecticide resistance in mosquito populations*.
- [3] Sharma, R., et al. (2020). *Biological control methods for mosquito management*.
- [4] Chandra, G., & Bhattacharjee, I. (2018). *Role of larvivorous fish in mosquito control*.
- [5] Becker, N. (2019). *Microbial control of mosquito larvae using Bacillus thuringiensis israelensis*.
- [6] Service, M. (2017). *Medical entomology for students*.
- [7] Centers for Disease Control and Prevention. (2022). *Integrated vector management strategies*.
- [8] Tiwari, S., et al. (2024). *Sustainable mosquito control approaches*.