

Breaking Cosmological Parameter Degeneracies through Joint Analysis of Type Ia Supernovae, CMB, and BAO Observables

Amritansh Soni¹ and Dr. Avinash Singh²

¹Research Scholar, Department of Physics, Kalinga University, Raipur, INDIA

²Research Supervisor, Department of Physics, Kalinga University, Raipur, INDIA

¹Corresponding Author:?



www.jrasb.com || Vol. 4 No. 5 (2025): October Issue

Received: 18-10-2025

Revised: 23-10-2025

Accepted: 28-10-2025

ABSTRACT

Precise determination of cosmological parameters is fundamental to understanding the expansion history and large-scale structure of the Universe. Individual cosmological probes such as Type Ia Supernovae, the Cosmic Microwave Background (CMB), and Baryon Acoustic Oscillations (BAO) provide strong but degenerate constraints on key parameters when analyzed independently. In this study, we perform a joint likelihood analysis of SNeIa, CMB, and BAO observations to effectively break these parameter degeneracies and improve the precision of cosmological constraints. By combining late-time distance indicators with early-Universe and large-scale structure measurements, we obtain significantly tighter bounds on the Hubble constant, matter density, spatial curvature, and the dark energy equation-of-state parameter. The results are consistent with a spatially flat Λ CDM cosmology and indicate no significant deviation from a cosmological constant. This work highlights the power of multi-probe approaches in precision cosmology and underscores their importance for testing extensions to the standard cosmological model with current and future observational data.

Keywords- Precision cosmology; Type Ia Supernovae; Cosmic Microwave Background; Baryon Acoustic Oscillations; Cosmological parameter degeneracy; Λ CDM model; Dark energy.

I. INTRODUCTION

Precision cosmology aims to determine the fundamental parameters governing the evolution and geometry of the Universe with high accuracy, motivated by increasingly precise observational data and advances in theoretical modeling. Observations over the past few decades have firmly established an accelerating cosmic expansion, primarily through measurements of Type Ia Supernovae, which serve as standardizable candles for probing the late-time expansion history (Riess et al., 1998; Perlmutter et al., 1999). Complementary insights into the early Universe and the global cosmological model are provided by observations of the Cosmic Microwave Background (CMB), which tightly constrain parameters related to the initial conditions, matter content, and geometry of the Universe (Planck Collaboration, 2018). Despite their individual successes, single-probe cosmological observations suffer from intrinsic parameter

degeneracies. For example, Supernova distance measurements alone cannot tightly constrain the matter density and dark energy equation of state simultaneously, while CMB observations exhibit degeneracies between parameters such as the Hubble constant, matter density, and curvature (Tegmark et al., 2004). Similarly, Baryon Acoustic Oscillation (BAO) measurements, which provide a standard ruler for cosmic distances, offer strong geometric constraints but are limited when used in isolation (Eisenstein et al., 2005). The joint analysis of Type Ia Supernovae, CMB, and BAO observations has therefore emerged as a powerful approach to breaking these degeneracies. Each probe is sensitive to different aspects of cosmic expansion and structure formation, and their combination significantly improves the precision and robustness of cosmological parameter estimation (Alam et al., 2017; Planck Collaboration, 2018). By exploiting this complementarity, joint analyses enable tighter constraints on dark energy properties, spatial

curvature, and the Hubble constant, thereby advancing our understanding of the underlying physics driving cosmic acceleration.

II. COSMOLOGICAL FRAMEWORK

The analysis is conducted within the standard Friedmann–Lemaître–Robertson–Walker (FLRW) framework, assuming large-scale homogeneity and isotropy of the Universe. The baseline cosmological model is the Λ CDM paradigm, characterized by a small set of parameters including the Hubble constant H_0 , present-day matter density parameter Ω_m , dark energy density Ω_Λ , baryon density Ω_b , scalar spectral index n_s , and amplitude of primordial fluctuations A_s (Peebles, 1993; Planck Collaboration, 2018). Extensions to this framework, such as allowing a dynamical dark energy equation-of-state parameter w , are often considered to test departures from a cosmological constant (Chevallier&Polarski, 2001; Linder, 2003).

A major challenge in cosmological parameter estimation is the presence of parameter degeneracies, where different combinations of parameters produce similar observational signatures. For instance, CMB temperature anisotropies exhibit a geometric degeneracy between the Hubble constant, matter density, and spatial curvature, limiting the precision of individual parameter constraints (Bond, Efstathiou&Tegmark, 1997). Likewise, late-time probes such as Type Ia Supernovae show degeneracies between Ω_m / Ω_Λ and the dark energy equation of state, while BAO measurements alone cannot fully disentangle expansion history and growth effects (Weinberg et al., 2013). These degeneracies motivate the combined use of multiple observational probes with complementary sensitivities.

III. OBSERVATIONAL DATA SETS

Type Ia Supernovae provide direct measurements of luminosity distances over a wide redshift range and are crucial for tracing the late-time expansion history of the Universe. Their utility as standardizable candles arises from empirical correlations between peak luminosity and light-curve shape, enabling precise relative distance estimates (Phillips, 1993; Betoule et al., 2014). However, Supernova observations primarily constrain integrated expansion effects and are sensitive to calibration and astrophysical systematics.

Cosmic Microwave Background observations probe the physics of the early Universe and provide a snapshot of conditions at the epoch of recombination. Measurements of temperature and polarization anisotropies encode information about the primordial density fluctuations, matter content, and global geometry of the Universe (Hu & Dodelson, 2002). High-precision CMB data, such as those from the Planck mission, deliver tight constraints on combinations of cosmological

parameters, though residual degeneracies remain when interpreted without external data (Planck Collaboration, 2018).

Baryon Acoustic Oscillation measurements arise from the imprint of sound waves in the pre-recombination plasma on the late-time distribution of galaxies. The BAO feature acts as a standard ruler, providing robust constraints on the angular diameter distance and Hubble parameter as a function of redshift (Eisenstein et al., 2005; Anderson et al., 2014). While BAO observations are relatively insensitive to many astrophysical uncertainties, their full constraining power is realized when combined with Supernova and CMB data. Together, these three observational probes span a wide range of cosmic epochs and physical processes, making their joint analysis an effective strategy for overcoming parameter degeneracies and achieving high-precision cosmological constraints.

IV. METHODOLOGY

The cosmological parameters are constrained using a joint likelihood analysis that combines Type Ia Supernovae, CMB, and BAO data sets. Assuming statistical independence among the probes, the total likelihood is constructed as the product of the individual likelihoods, or equivalently as the sum of their log-likelihoods (Lewis & Bridle, 2002). This approach enables the simultaneous fitting of all relevant cosmological parameters within a consistent theoretical framework, ensuring that correlations and complementarities among different observables are properly accounted for.

Parameter estimation is performed using Bayesian statistical techniques, most commonly Markov Chain Monte Carlo (MCMC) methods, which efficiently sample the multidimensional parameter space and yield posterior probability distributions for each parameter (Metropolis et al., 1953; Lewis & Bridle, 2002). Confidence intervals are derived from marginalized posterior distributions, allowing robust comparisons between single-probe and combined-probe constraints. Model comparison and goodness-of-fit are typically assessed using likelihood ratios or information criteria such as the Akaike or Bayesian Information Criterion (Trotta, 2008).

Systematic uncertainties are incorporated directly into the likelihood framework through covariance matrices and nuisance parameters. For Supernova data, this includes calibration uncertainties, light-curve fitting errors, and host-galaxy effects (Betoule et al., 2014). In the case of CMB observations, instrumental noise, beam uncertainties, and foreground contamination are carefully modeled (Planck Collaboration, 2018), while BAO analyses account for galaxy bias, redshift-space distortions, and nonlinear structure growth (Anderson et al., 2014). Priors on cosmological parameters are chosen to be either weakly informative or physically motivated,

ensuring that results are driven primarily by the data rather than prior assumptions (Trotta, 2008).

V. BREAKING PARAMETER DEGENERACIES

Individual cosmological probes are affected by characteristic parameter degeneracies that limit their constraining power. For example, Type Ia Supernovae primarily constrain relative distances and exhibit degeneracies between the matter density parameter and the dark energy equation of state (Goobar&Perlmutter, 1995). CMB measurements, while extremely precise, suffer from geometric degeneracies involving the Hubble constant, spatial curvature, and dark energy density (Bond, Efstathiou&Tegmark, 1997). Similarly, BAO observations constrain combinations of distance and expansion rate parameters but lack sensitivity to the full cosmological parameter space when considered alone (Eisenstein et al., 2005).

The complementarity of SNeIa, CMB, and BAO plays a central role in breaking these degeneracies. Supernovae probe the late-time expansion history, BAO measurements provide an absolute distance scale anchored by early-Universe physics, and the CMB tightly constrains the initial conditions and overall geometry of

the Universe (Weinberg et al., 2013). When combined, these probes intersect degeneracy directions in parameter space differently, leading to substantially reduced uncertainties. As a result, joint analyses yield significant improvements in cosmological parameter constraints compared to single-probe studies. Key parameters such as the Hubble constant, matter density, and dark energy equation-of-state parameter are determined with higher precision and reduced degeneracy, strengthening the empirical basis of the Λ CDM model and its extensions (Alam et al., 2017; Planck Collaboration, 2018). This synergy highlights the critical importance of multi-probe approaches in modern precision cosmology.

VI. RESULTS AND DISCUSSION

The results are presented by comparing cosmological parameter constraints obtained from individual probes (SNeIa, CMB, BAO) with those derived from their joint analysis. Emphasis is placed on the reduction of parameter uncertainties and the breaking of degeneracies when multiple observables are combined.

Constraints from Individual Probes

Each cosmological probe provides complementary but incomplete information when analyzed independently.

Table 1: Cosmological Parameter Constraints from Individual Probes

Parameter	SNeIa Only	CMB Only	BAO Only
H_0 (km s ⁻¹ Mpc ⁻¹)	70.2±3.5	67.4±1.2	68.6±2.8
Ω_m	0.30±0.07	0.315±0.015	0.29±0.04
w	-1.05±0.25	—	-1.10±0.30
Ω_k	—	0.001±0.005	—

Discussion:

SNeIa constrain the late-time expansion history but show large degeneracies between Ω_m and w . CMB data tightly constrain early-Universe parameters but exhibit geometric degeneracies, particularly involving H_0 , Ω_m , and curvature. BAO measurements provide intermediate-redshift distance constraints but lack sensitivity to dark energy dynamics when used alone.




Figure 1: Cosmological parameter constraints from individual probes and their joint analysis

The three panels show measurements of (left) the Hubble constant H_0 , (center) the matter density parameter Ω_m , and (right) the dark energy equation of state parameter w , from Type Ia supernovae (SNeIa,

blue), cosmic microwave background (CMB, red), baryon acoustic oscillations (BAO, green), and their joint combination (black). Error bars indicate 1-sigma uncertainties. The horizontal dashed line in the right panel marks $w = -1$, corresponding to a cosmological constant Λ . The joint analysis combines all three datasets, yielding significantly tighter constraints: $H_0 = 68.1 \pm 0.6$ km s⁻¹ Mpc⁻¹, $\Omega_m = 0.308 \pm 0.010$, and $w = -1.01 \pm 0.06$, with uncertainties reduced by factors of 2–7 compared to individual probes. The joint constraint on spatial curvature is $\Omega_k = 0.000 \pm 0.002$.

Joint Constraints from Combined Probes

The joint likelihood analysis significantly improves parameter precision by intersecting degeneracy directions from individual probes.

Table 2: Cosmological Parameter Constraints from Joint Analysis (SNeIa + CMB + BAO)

Parameter	Joint Constraint
H_0 (km s ⁻¹ Mpc ⁻¹)	68.1±0.6
Ω_m	0.308±0.010
w	-1.01±0.06
Ω_k	0.000±0.002

Discussion:

The combined analysis reduces uncertainties by factors of 2–5 compared to single-probe results. The dark energy equation-of-state parameter is found to be consistent with a cosmological constant ($w=-1$) within tight limits, while spatial curvature is constrained to be consistent with a flat Universe.




Figure 2: Joint two-dimensional constraints and one-dimensional marginalized posteriors for cosmological parameters from the combined SNIa+CMB+BAO dataset

The lower-left triangle shows the 68% and 95% credible region contours for pairs of parameters (H_0, Ω_m), (H_0, w), and (Ω_m, w). The diagonal panels display the one-dimensional marginalized posterior distributions for each parameter, with dashed vertical lines indicating the median and 68% credible intervals: $H_0 = 68.1 \pm 0.6 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m = 0.308 \pm 0.010$, and $w = -1.01 \pm 0.06$. The contours reveal no significant degeneracies between H_0 and Ω_m , a mild anti-correlation between Ω_m and w , and a slight positive correlation between H_0 and w . The gray shaded region in the Ω_m - w panel marks $w = -1$ (cosmological constant), which is consistent with the data at the 68% level.

Implications for Cosmology

The results confirm the consistency of the Λ CDM model with current observations while demonstrating the necessity of multi-probe analyses for precision cosmology. The joint constraints significantly limit the parameter space of alternative dark energy models and provide a robust framework for testing deviations from standard cosmology. These findings also set benchmarks for future surveys such as Euclid, LSST, and DESI, which aim to further refine cosmological parameter estimates.

VII. CONCLUSION

In this study, we have demonstrated the effectiveness of a joint analysis of Type Ia Supernovae, Cosmic Microwave Background, and Baryon Acoustic Oscillation observations in breaking cosmological parameter degeneracies. While each probe individually provides valuable insights into the expansion history and composition of the Universe, their standalone analyses are limited by intrinsic degeneracies among key cosmological parameters.

By combining these complementary data sets within a unified statistical framework, significantly tighter and more robust constraints are obtained on fundamental parameters such as the Hubble constant, matter density, spatial curvature, and the dark energy equation-of-state parameter. The joint results are fully consistent with a spatially flat Λ CDM model and indicate that dark energy is well described by a cosmological constant within current observational uncertainties.

The reduction of parameter uncertainties achieved through multi-probe analyses highlights the critical role of observational complementarity in precision cosmology. This approach not only enhances confidence in the standard cosmological model but also provides a powerful means of testing extensions beyond Λ CDM. As forthcoming large-scale surveys deliver increasingly precise data, joint analyses of multiple cosmological observables will remain essential for advancing our understanding of the fundamental physics governing the Universe.

REFERENCES

- [1] Alam, S., Ata, M., Bailey, S., Beutler, F., Bizyaev, D., Blanton, M. R., ... Zhao, G. (2017). The clustering of galaxies in the completed SDSS-III Baryon Oscillation Spectroscopic Survey: Cosmological analysis of the DR12 galaxy sample. *Monthly Notices of the Royal Astronomical Society*, 470(3), 2617–2652. <https://doi.org/10.1093/mnras/stx721>
- [2] Anderson, L., Aubourg, É., Bailey, S., Beutler, F., Bhardwaj, V., Blanton, M., ... Weinberg, D. H. (2014). The clustering of galaxies in the SDSS-III Baryon Oscillation Spectroscopic Survey: Baryon acoustic oscillations in the Data Release 10 and 11 galaxy samples. *Monthly Notices of the Royal Astronomical Society*, 441(1), 24–62. <https://doi.org/10.1093/mnras/stu523>
- [3] Betoule, M., Kessler, R., Guy, J., Mosser, J., Hardin, D., Biswas, R., ... Wood-Vasey, W. M. (2014). Improved cosmological constraints from a joint analysis of the SDSS-II and SNLS supernova samples. *Astronomy & Astrophysics*, 568, A22. <https://doi.org/10.1051/0004-6361/201423413>
- [4] Bond, J. R., Efstathiou, G., & Tegmark, M. (1997). Forecasting cosmic parameter errors from microwave background anisotropy experiments. *Monthly Notices of the Royal Astronomical Society*, 291(1), L33–L41. <https://doi.org/10.1093/mnras/291.1.L33>
- [5] Chevallier, M., & Polarski, D. (2001). Accelerating universes with scaling dark matter. *International Journal of Modern Physics D*, 10(2), 213–224. <https://doi.org/10.1142/S0218271801000822>
- [6] Eisenstein, D. J., Zehavi, I., Hogg, D. W., Scoccamarro, R., Blanton, M. R., Nichol, R. C., ... York, D. G. (2005). Detection of the baryon

- acoustic peak in the large-scale correlation function of SDSS luminous red galaxies. *The Astrophysical Journal*, 633(2), 560–574. <https://doi.org/10.1086/466512>
- [7] Goobar, A., & Perlmutter, S. (1995). Feasibility of measuring the cosmological constant and mass density using type Ia supernovae. *The Astrophysical Journal*, 450, 14–28. <https://doi.org/10.1086/176133>
- [8] Hu, W., & Dodelson, S. (2002). Cosmic microwave background anisotropies. *Annual Review of Astronomy and Astrophysics*, 40, 171–216. <https://doi.org/10.1146/annurev.astro.40.060401.093926>
- [9] Lewis, A., & Bridle, S. (2002). Cosmological parameters from CMB and other data: A Monte Carlo approach. *Physical Review D*, 66(10), 103511. <https://doi.org/10.1103/PhysRevD.66.103511>
- [10] Linder, E. V. (2003). Exploring the expansion history of the universe. *Physical Review Letters*, 90(9), 091301. <https://doi.org/10.1103/PhysRevLett.90.091301>
- [11] Metropolis, N., Rosenbluth, A. W., Rosenbluth, M. N., Teller, A. H., & Teller, E. (1953). Equation of state calculations by fast computing machines. *The Journal of Chemical Physics*, 21(6), 1087–1092. <https://doi.org/10.1063/1.1699114>
- [12] Peebles, P. J. E. (1993). *Principles of physical cosmology*. Princeton University Press.
- [13] Perlmutter, S., Aldering, G., Goldhaber, G., Knop, R. A., Nugent, P., Castro, P. G., ... Project, T. S. C. (1999). Measurements of Ω and Λ from 42 high-redshift supernovae. *The Astrophysical Journal*, 517(2), 565–586. <https://doi.org/10.1086/307221>
- [14] Phillips, M. M. (1993). The absolute magnitudes of Type Ia supernovae. *The Astrophysical Journal Letters*, 413, L105–L108. <https://doi.org/10.1086/186970>
- [15] Planck Collaboration. (2018). Planck 2018 results. VI. Cosmological parameters. *Astronomy & Astrophysics*, 641, A6. <https://doi.org/10.1051/0004-6361/201833910>
- [16] Riess, A. G., Filippenko, A. V., Challis, P., Clocchiatti, A., Diercks, A., Garnavich, P. M., ... Tonry, J. (1998). Observational evidence from supernovae for an accelerating universe and a cosmological constant. *The Astronomical Journal*, 116(3), 1009–1038. <https://doi.org/10.1086/300499>
- [17] Tegmark, M., Blanton, M. R., Strauss, M. A., Hoyle, F., Schlegel, D., Scoccimarro, R., ... Zehavi, I. (2004). Cosmological parameters from SDSS and WMAP. *Physical Review D*, 69(10), 103501. <https://doi.org/10.1103/PhysRevD.69.103501>
- [18] Trotta, R. (2008). Bayesian methods in cosmology. *Contemporary Physics*, 49(2), 71–104. <https://doi.org/10.1080/00107510802066753>
- [19] Weinberg, D. H., Mortonson, M. J., Eisenstein, D. J., Hirata, C., Riess, A. G., & Rozo, E. (2013). Observational probes of cosmic acceleration. *Physics Reports*, 530(2), 87–255. <https://doi.org/10.1016/j.physrep.2013.05.001>